

Homework Set 10

Due October 30

- Construct a spherical tensor of rank 1 out of two different vectors $\mathbf{U} = (U_x, U_y, U_z)$ and $\mathbf{V} = (V_x, V_y, V_z)$.
Explicitly write $T_{\pm 1,0}^{(1)}$ in terms of $U_{x,y,z}$ and $V_{x,y,z}$.
 - Construct a spherical tensor of rank 2 out of two different vectors \mathbf{U} and \mathbf{V} .
Write down explicitly $T_{\pm 2,\pm 1,0}^{(2)}$ in terms of $U_{x,y,z}$ and $V_{x,y,z}$.
- Consider a spinless particle bound to a fixed center by a central force potential.

- Relate, as much as possible, the matrix elements

$$\langle n', l', m' | \mp \frac{1}{\sqrt{2}}(x \pm iy) | n, l, m \rangle \quad (1)$$

and

$$\langle n', l', m' | z | n, l, m \rangle \quad (2)$$

using *only* the Wigner-Eckart theorem. Make sure to state under what conditions the matrix elements are nonvanishing.

- Do the same problem using wave functions

$$\psi(\mathbf{x}) = R_{nl}(r)Y_l^m(\theta, \phi) \quad (3)$$

- Write xy , xz , and $(x^2 - y^2)$ as components of a spherical (irreducible) tensor of rank 2.
 - The expectation value

$$Q \equiv e \langle \alpha, j, m = j | (3z^2 - r^2) | \alpha, j, m = j \rangle \quad (4)$$

is known as the *quadrupole moment*. Evaluate

$$e \langle \alpha, j, m' | (x^2 - y^2) | \alpha, j, m = j \rangle \quad (5)$$

where $m' = j, j-1, j-2, \dots$, in terms of Q and appropriate Clebsch-Gordan coefficients